Creative Enzymes®

β-1,3-Galactosyl-O-glycosyl-Glykoprotein β-1,3-N-Acetylglucosaminyltransferase

Cat. No. EXWM-2372

Lot. No. (See product label)

Einleitung

Beschreibung	vgl. EC 2.4.1.102 (β-1,3-galactosyl-O-glycosyl-glycoprotein β-1,6-N-acetylglucosaminyltransferase), EC 2.4.1.147 (acetylgalactosaminyl-O-glycosyl-glycoprotein β-1,3-N-acetylglucosaminyltransferase) und EC 2.4.1.148 (acetylgalactosaminyl-O-glycosyl-glycoprotein β-1,6-N-acetylglucosaminyltransferase).
Synonyme	O-glycosyl-oligosaccharid-glykoprotein N-Acetylglucosaminyltransferase II; Uridin-Diphosphoacetylglucosamin-Mucin β(1→3)-Acetylglucosaminyltransferase (verlängernd); Verlängerung 3β-GalNAc-Transferase; UDP-N-Acetyl-D-glucosamin:O-glycosyl-Glykoprotein (N-Acetyl-D-glucosamin zu β-D-Galactose von β-D-Galactosyl-1,3-(N-Acetyl-D-glucosaminyl-1,6)-N-Acetyl-D-galactosaminyl-R) β-1,3-N-Acetyl-D-glucosaminyltransferase

Produktinformation

Form	Flüssigkeit oder lyophilisiertes Pulver
EC-Nummer	EC 2.4.1.146
CAS-Nummer	87927-99-9
Reaktion	UDP-N-acetyl-D-glucosamin + β-D-galactosyl-(1→3)-[N-acetyl-D-glucosaminyl-(1→6)]-N-acetyl-D-galactosaminyl-R = UDP + N-acetyl-β-D-glucosaminyl-(1→3)-β-D-galactosyl-(1→3)-[N-acetyl-β-D-glucosaminyl-(1→6)]-N-acetyl-D-galactosaminyl-R
Hinweise	Dieser Artikel erfordert eine maßgeschneiderte Produktion, und die Lieferzeit beträgt zwischen 5 und 9 Wochen. Wir können nach Ihren Spezifikationen maßgeschneidert produzieren.

Lager- und Versandinformation

Lagerung	Lagern Sie es kurzfristig bei +4 ℃. Für die Langzeitlagerung lagern Sie es bei -20 ℃~-80 ℃.