Creative Enzymes®

mannosyl-Glycoprotein Endo-β-N-Acetylglucosaminidase

Cat. No. EXWM-3959

Lot. No. (See product label)

Einleitung

Beschreibung	Eine Gruppe verwandter Enzyme.
Synonyme	N,N'-diacetylchitobiosyl β-N-acetylglucosaminidase; endo-β-N-acetylglucosaminidase; mannosyl-glycoprotein endo-β-N-acetylglucosamidase; di-N-acetylchitobiosyl β-N-acetylglucosaminidase; endo-β-acetylglucosaminidase; endo-β-(1→4)-N-acetylglucosaminidase; mannosyl-glycoprotein 1,4-N-acetamidodeoxy-β-D-glycohydrolase; endoglycosidase S; endo-N-acetyl-β-D-glucosaminidase; endo-N-acetyl-β-glucosaminidase; endo-β-N-acetylglucosaminidase D; endo-β-N-acetylglucosaminidase F; endo-β-N-acetylglucosaminidase H; endo-β-N-acetylglucosaminidase L; glycopeptide-D-mannosyl-4-N-(N-acetyl-D-glucosaminyl)2-asparagine 1,4-N-acetyl-β-glucosaminohydrolase; endoglycosidase H

Produktinformation

Form	Flüssigkeit oder lyophilisiertes Pulver
EC-Nummer	EC 3.2.1.96
CAS-Nummer	37278-88-9
Reaktion	Endohydrolyse der N,N'-diacetylchitobiosyl-Einheit in hochmannosereichen Glykoproteinen und Glykpeptiden, die die -[Man(GlcNAc)2]Asn- Struktur enthalten. Ein N-Acetyl-D-Glucosamin-Rest bleibt an dem Protein gebunden; der Rest des Oligosaccharids wird intakt freigesetzt.
Hinweise	Dieser Artikel erfordert eine maßgeschneiderte Produktion, und die Lieferzeit beträgt zwischen 5 und 9 Wochen. Wir können nach Ihren Spezifikationen maßgeschneidert produzieren.

Lager- und Versandinformation

Lagerung	Lagern Sie es kurzfristig bei +4 ℃. Für die Langzeitlagerung lagern Sie es bei -20 ℃~-80 ℃.